

Ready Rumen



Rapid growth and convenience

Suitable for calves reared for both beef calves and dairy heifer replacements.

NRM Ready Rumen is a measured blend of high protein pellets, straw and molasses. Ready Rumen also contains Bovatec® for coccidiosis control. The roughage included is chopped to length and supplied at the optimum ratio to develop the rumen as early as possible in young calves. Ready Rumen is a cost effective, hassle free feed that requires no added roughage, vitamins or minerals – ideal for the once-a-day feeding system.

Feeding Recommendation

Introduce Ready Rumen gradually from birth and feed ad-lib through to 84 days of age. On this system, calves do not need access to pasture until 8 - 10 weeks.

Always allow free access to fresh, clean water.

Key Benefits & Features

- Moozlee flavourants for consistent palatability and early feed intakes.
- Includes a high protein pellet.
- Balanced energy/protein to maximise growth.
- Convenience of measured straw intake for rumen development.
- Stabilises rumen environment to reduce risk of acidosis resulting from high concentrate intake.
- Bovatec® for coccidiosis protection.
- Recyclable packaging.
- PKE free.



Feeding Rates

AGE	NRM POWER WHEY*	NRM READY RUMEN	TARGET WEIGHT
Day 0 - 4	colostrum	ad-lib	40kg Body Weight min
Day 5 - 10	2L twice-a-day	ad-lib	
Day 11 - 21	2.5L twice-a-day	ad-lib	
Day 22 - wean	3L twice-a-day	ad-lib	28 days = 50kg
Day 50 - 66		ad-lib	66 days = 75kg
Day 67 - 84		ad-lib	84 days = 100kg

* 125 - 150g of NRM Power Whey with water to make up 1L

Storage

Please ensure product is stored in a cool, dry and vermin free environment.

Caution

Do not feed to any other animal species other than those stipulated on the label.

Ingredients

Selected from: Grain and grain by-products, various plant proteins, food by-products, lucerne chaff, minerals, vegetable oils, mould inhibitor, vitamins and trace minerals, Bovatec® 20CC.

Feed To Grow

Typical Analysis

(approximate on a DM basis)

Protein	19.5%
Moisture	12.5%



Ready Rumen



Important

Bovatec® is registered pursuant to the AVCM Act 1997, No. A009679.

See www.nzfsa.govt.nz/acvm for Bovatec® registration conditions.

Ensure young animals are adopted to medicated feed two - three weeks prior to weaning, as coccidial challenge is most likely in the eight weeks after weaning. In any case initiate treatment at least two days prior to expected coccidial challenge.

Feeding Recommendation for Coccidiosis Control

For coccidiosis control, minimum dose rate of 5mg Bovatec®/kg liveweight (1mg lasalocid sodium/kg liveweight) is required per day. Up to twice this dose rate is indicated where severe coccidial challenge is expected or a wide range of individual liveweights occur within a group.

Feeding Rates for Coccidiosis Control

CALF WEIGHT (LIVEWEIGHT)	FEED PER DAY
40kg	0.4kg
50kg	0.5kg
60kg	0.6kg
70kg	0.7kg
80kg	0.8kg
90kg	0.9kg
100kg	1.0kg
110kg	1.1kg

Calf feed with Bovatec® 20CC, ACVM Registration Number A009679


BOVATEC® FOR ANIMAL TREATMENT ONLY
Contains lasalocid sodium at 99mg/kg of feed

Indications: Coccidiosis control in cattle, calves and goats.

Contraindications: Care must be exercised when feeding concurrently with other antimicrobials. Do not feed with other ionophores, e.g. monensin capsules, liquid or premix. Do not exceed recommended dose rates. Do not feed more than a total of 400mg lasalocid/head/day to lactating cattle. Not to be used for single-dose treatment. Do not allow dogs, horses (or other equids), alpacas (or other camelids) access to feeds containing Bovatec, as ingestion by these species may be fatal.

Withholding period: Meat: nil Milk: cattle - nil, goats - 35 days

Feed To Grow



Feed To Grow 0800 800 380 www.nrm.co.nz

THE QUALITY OF NRM PRODUCTS IS GUARANTEED

We guarantee the quality of our NRM feeds and animal health products. If you have any queries please contact us.

