

Calf Milk Finisher with Bovatec®

Complete milk replacer for young calves

An economical calf milk replacer, formulated as a full replacement of natural cow's milk which can be fed to calves after the initial colostrum period.

Directions for use

Decide on which feeding option from the tables below best applies to your situation.

Mixing ratio: Typically 150g NRM Calf Milk Finisher made up to one litre with water when fed twice daily, 300g NRM Calf Milk Finisher to made up to one litre of water when fed once per day.

Dissolve NRM Calf Milk Finisher powder in hot (not boiling) water. Mix to a creamy, smooth consistency and add cool water to make up the required amount. Feed at body temperature (38-40°C). Mix only when required.

Feed after a good intake of colostrum has been achieved (from four days after birth). Keep buckets, bottles and other equipment clean.

Ensure calves have access to long fibre and quality calf feed.



PROTEIN



MINERALS & VITAMINS



COCCIDIOSTAT

Feeding Programme

ONCE A DAY SYSTEM

AGE IN DAYS	NO. FEEDS/DAY	LITRES/FEED	CONCENTRATION
1-4	2	2.5	Colostrum Only
5-14	2	2	150g/litre
15-28	1	2.5	300g/litre
29-weaning	1	2.5	300g/litre

TWICE A DAY SYSTEM

AGE IN DAYS	NO. FEEDS/DAY	LITRES/FEED	CONCENTRATION
1-4	2	2.5	Colostrum Only
5-14	2	2	150g/litre
15-28	2	2.5	150g/litre
29-weaning	2	2.5	150g/litre

Ingredients selected from

Whole milk powder, whey powder, vegetable fats, soy protein, vitamins, minerals, Bovatec® 20CC, Bovatec Technical. Additional copper, Bovatec Technical or selenium should only be given following consultation with a nutritionist or veterinarian.

Storage

Store between 10 and 20°C in clean, dry conditions. Do not expose to direct sunlight and avoid contact with floor and walls.

Caution: Do not feed to any animal species other than those stipulated on the label.

Calf Milk Finisher with Bovatec®

Complete milk replacer for young calves

Typical analysis (approximate on an as fed basis)

Crude Protein	24%
Crude Fat	20%
Lactose	38%

Calf Milk Finisher with Bovatec Technical®

FOR ANIMAL TREATMENT ONLY

CONTAINS: Bovatec Technical at 111mg/kg (100mg/kg of Lasalocid).

INDICATIONS: As an aid in the control of coccidiosis caused by *Eimeria* species in goats and cattle. Feed a 50kg calf 0.5kg/day of NRM Calf Milk Finisher with Bovatec to provide 1mg lasalocid/kg body weight. Up to 2mg lasalocid/kg live weight is indicated where severe coccidial challenge is expected or where a wide range of individual live weights occur within a group.

CONTRAINDICATIONS: Care must be exercised when feeding concurrently with other antimicrobials. Do not feed with other ionophores, e.g. monensin capsules, liquid or premix. Do not exceed the recommended dose rates. Not to be used for single-dose treatment.

CAUTION: Do not feed to bobby calves. Do not allow dogs, horses (or other equids), alpacas (or other camelids) access to feeds containing Bovatec®, as ingestion by these species may be fatal. This product must not be used for growth promotion in ruminants intended for human consumption.

WITHHOLDING PERIOD

It is an offence for users of this product to cause residues exceeding the relevant MRL (Maximum Residue Limits of Agricultural Compounds) in the New Zealand Food Standard.

Meat: Cattle, Nil

Milk: Cattle, Nil

Bovatec Technical® is registered pursuant to the ACVM Act, 1997. ACVM Registration Number A10829.

See www.nzfsa.govt.nz/acvm for registration conditions.

By law the user must take due care, obtaining expert advice if necessary to avoid unnecessary pain and distress when using the product other than as directed on the label.

THE QUALITY OF NRM PRODUCTS IS GUARANTEED

If you have any queries, please contact us.

0800 800 380
www.nrm.co.nz