

Calf Milk Replacer with Bovatec®

Whole milk replacer for young calves

An all milk powder blend providing excellent calf response to the natural food sources of milk fat, milk protein and sugar.

Directions for use

Decide on which feeding option from the tables below best applies to your situation.

Mixing ratio: Typically 125g Calf Milk Replacer made up to one litre with water when fed twice daily, 250g Calf Milk Replacer made up to one litre with water when fed once daily.

Dissolve Calf Milk Replacer powder in hot (not boiling) water. Mix to a creamy, smooth consistency and add cool water to make up the required amount. Feed at body temperature (38-40°C). Mix only when required.

Feed after a good intake of colostrum has been achieved (from four days after birth). Keep buckets, bottles and other equipment clean.

Ensure calves have access to long fibre and quality calf feed.



MILK FAT



MILK PROTEIN



COCCIDIOSTAT

Feeding Programme

ONCE A DAY SYSTEM

AGE IN DAYS	NO. FEEDS/DAY	LITRES/FEED	CONCENTRATION	DAILY INTAKE OF PELLETS/MUESLI (g)
1-4	2	2.5	Colostrum Only	-
5-14	2	2	125g/litre	50-100
15-28	1	2.5	250g/litre	100-350
29-weaning	1	2.5	250g/litre	350-1000

TWICE A DAY SYSTEM

AGE IN DAYS	NO. FEEDS/DAY	LITRES/FEED	CONCENTRATION	DAILY INTAKE OF PELLETS/MUESLI (g)
1-4	2	2.5	Colostrum Only	-
5-14	2	2	125g/litre	50-100
15-28	2	2.5	125g/litre	100-350
29-weaning	2	2.5	125g/litre	350-1000

Ingredients selected from

Milk Powder, Vitamin A, D3, E, K3, B1, B2, B3, B5, B6, B12, Ascorbic Acid, Potassium, Lasalocid (Bovatec), Niacin, Pantothenic Acid, Biotin, Cobalt (organic), Copper (organic), Dextrose, Ferrous Iron, Iodine, Selenium (organic), Magnesium, Manganese (organic), Zinc (organic), Antioxidants, Lecithin. Additional copper or selenium should only be given following consultation with a nutritionist or veterinarian.

Storage

Store between 10 and 20°C in clean, dry conditions. Do not expose to direct sunlight and avoid contact with floor and walls.

Caution: Do not feed to any animal species other than those stipulated on the label.

Calf Milk Replacer with Bovatec®

Whole milk replacer for young calves

Typical analysis (approximate on an as fed basis)

Crude Protein	25%
Crude Fat	20%
Lactose	44%
Ash	6%

Calf Milk Replacer with Bovatec Technical®

FOR ANIMAL TREATMENT ONLY

CONTAINS: Bovatec Technical at 111mg/kg (100mg/kg of Lasalocid).

INDICATIONS: As an aid in the control of coccidiosis caused by *Eimeria* species in goats and cattle. Feed a 50kg calf 0.5kg/day of NRM Calf Milk Finisher with Bovatec to provide 1mg lasalocid/kg body weight. Up to 2mg lasalocid/kg live weight is indicated where severe coccidial challenge is expected or where a wide range of individual live weights occur within a group.

CONTRAINDICATIONS: Care must be exercised when feeding concurrently with other antimicrobials. Do not feed with other ionophores, e.g. monensin capsules, liquid or premix. Do not exceed the recommended dose rates. Not to be used for single-dose treatment.

CAUTION: Do not feed to bobby calves. Do not allow dogs, horses (or other equids), alpacas (or other camelids) access to feeds containing Bovatec®, as ingestion by these species may be fatal. This product must not be used for growth promotion in ruminants intended for human consumption.

WITHHOLDING PERIOD

It is an offence for users of this product to cause residues exceeding the relevant MRL (Maximum Residue Limits of Agricultural Compounds) in the New Zealand Food Standard.

Meat: Cattle, Nil

Milk: Cattle, Nil

Bovatec Technical® is registered pursuant to the ACVM Act, 1997. ACVM Registration Number A10829. See www.nzfsa.govt.nz/acvm for registration conditions.

By law the user must take due care, obtaining expert advice if necessary to avoid unnecessary pain and distress when using the product other than as directed on the label.