



Chick Starter Crumble

with Coccidiostat FOR ANIMAL TREATMENT ONLY

A complete feed for young chickens providing nutrients essential for lean growth and skeletal development

NRM Chick Starter Crumble is a balanced feed designed to promote early bone, frame and feather development, rapid muscle deposition and a high health status in young chickens. Contains a proven coccidiostat to help protect young chicks against coccidiosis until they build up natural immunity.

KEY BENEFITS AND FEATURES

NRM Chick Starter Crumble contains:

- Essential nutrients locked into every feed pellet, which when crumbled, improves appeal and accessibility for young chicks.
- Elevated levels of protein balanced according to known amino acid requirements to support growth.
- Additives including plant extracts to support gut health, especially useful in chicks when young and vulnerable.
- Major minerals, trace elements, and vitamins to support skeletal growth, muscle development, and general health of chicks, and aids peace of mind for owners too.
- A proven coccidiostat to aid against protozoal parasites that cause coccidiosis a debilitating disease which can reduce growth rates and even kill young birds.
- Enzymes to improve the digestion of grains for improved feed utilisation.




Product available in 10kg or 20kg.









Approximate weeks depending on breed and season







Chick Starter Crumble

with Coccidiostat
FOR ANIMAL TREATMENT ONLY

FEEDING RECOMMENDATIONS

Feed NRM Chick Starter Crumble to appetite from one day old for up to eight weeks. Clean and replenish feed containers regularly to prevent the accumulation of stale or contaminated feed.

Always ensure birds have access to clean, fresh water. Not recommended for gamebirds, ducklings and other waterfowl.

For chick rearing advice and alternative product suggestions for waterfowl and game birds, please use the OR code below.

RECYCLE ME

NRM bags with the Recycle Me logo use 25% less plastic per bag compared to previous versions, giving them a lower carbon footprint and enable their recyclability through Agrecovery.

Scan QR below for more information.

TYPICAL ANALYSIS

Nutrient	As Fed Basis	Dry Matter Basis
Dry Matter %	88	100
Crude Protein %	19	21.5
Available Lysine g/kg	9.5	10.8

Variations in nutritional values may occur due to natural variability in feed ingredients. Our FeedSafe NZ certified sites have well developed systems to minimise any variation from the typical levels shown.

INGREDIENTS SELECTED FROM

Grain and grain by-products, plant proteins, vegetable oils, enzymes, amino acids, limestone, mono or dicalcium phosphate, salt, sodium bicarbonate, vitamins, trace minerals, organic acids and plant extracts, and coccidiostat (Bovatec® 20CC).

May contain or be sourced from genetically modified source crops.

STORAGE

Store in a cool, dry and vermin free environment out of direct sunlight.

Do not feed crumble which has become wet and moulded.

WARNING STATEMENTS

KEEP OUT OF THE REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ THE LABEL BEFORE USE

NRM Chick Starter Crumble with 450g Bovatec® 20CC/tonne.

ACTIVE INGREDIENT: Lasalocid sodium 90mg per kilogram.

CONTAINS: 450g Bovatec® 20CC per tonne, which is registered pursuant to the ACVM Act 1997, No. A009679. See www.foodsafety.govt.nz for registration conditions.

DIRECTIONS FOR USE: This product must not be used for growth promotion in ruminants intended for human consumption. By law the user must take due care, obtaining expert advice if necessary to avoid unnecessary pain and distress when using the product other than as directed on the label.

INDICATIONS: Delivers 90mg Lasalocid sodium per kg as 450g Bovatec® 20CC/ tonne, as an aid in the prevention of coccidiosis caused by *Eimeria* species in broiler chickens, replacement layer and breeder pullets on deep litter or range.

DOSE RATE: As an aid in the prevention of coccidiosis, feed continuously NRM Chick Starter Crumble fresh daily to appetite

as the only ration. Discontinue use 14 DAYS before egg laying commences.

CONTRAINDICATIONS: Care must be exercised when feeding concurrently with other antimicrobials. Do not feed with other ionophores, e.g., monensin capsules, liquid, or pre-mix. Do not exceed recommended dose rates. Not to be used for single dose treatment.

CAUTION: Do not allow dogs, horses (or other equids), alpacas (or other camelids) access to feeds containing Bovatec® 20CC, as ingestion by these species may be fatal.

WITHHOLDING PERIODS: It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Poultry:

- Meat: Nil
- Eggs: Eggs from treated birds must not be sold for human consumption for 14 days following the last treatment.

DISPOSAL: : Avoid any contamination of any water supply with this feed or packaging.

DO NOT FEED TO ANIMALS OTHER THAN THOSE STIPULATED ON THIS LABEL $\,$





